

Dimension-six Proton Decays in the Modified Missing Doublet SU(5) Model

J. Hisano^{a)}, Yasunori Nomura^{b)}, and T. Yanagida^{b)}

a) *Theory Group, KEK, Tsukuba, Ibaraki 305, Japan*

b) *Department of Physics, University of Tokyo, Tokyo 113, Japan*

Abstract

Dimension-five operators for nucleon decays are suppressed in the modified missing doublet (MMD) model in the supersymmetric SU(5) grand unification. We show that nonrenormalizable interactions decrease the unification scale in the MMD model which increases the nucleon decay rate of dimension-six operators by a significant amount. We find that the theoretical lower bound on the proton life time $\tau(p \rightarrow e^+ \pi^0)$ is within the observable range at SuperKamiokande.

The doublet-triplet splitting is one of the most serious problems in the supersymmetric (SUSY) grand unified theory (GUT). Various interesting resolutions to this difficult problem have been proposed [1][2][3][4][5][6]. Among them the modified missing doublet (MMD) SU(5) model [2] is the most attractive one, since it explains very naturally the low value of the SU(3)_C gauge coupling constant $\alpha_3(m_Z) = 0.118 \pm 0.004$ [7] by GUT-scale threshold corrections [8]. Furthermore, the dangerous dimension-five operators for nucleon decays [9] are strongly suppressed [2] owing to an additional chiral symmetry in the MMD model.

If the nucleon decay due to the dimension-five operators are outside the reach at SuperKamiokande, the dimension-six nucleon decays such as $p \rightarrow e^+ \pi^0$ are clearly important to test the GUT. In this letter, we point out that introduction of non-renormalizable operators may decrease the GUT scale by a factor about 3 for a reasonable parameter region in the MMD SU(5) model. As a consequence the nucleon decays by the GUT gauge-boson exchanges become ~ 100 times faster than the previous estimate [10]. The present analysis, therefore, suggests that the theoretical lower bound for proton decay of the dominant channel $p \rightarrow e^+ \pi^0$ is within the observable range at the SuperKamiokande detector.¹

The original missing doublet (MD) model [1] in the SUSY SU(5) GUT consists of the following chiral supermultiplets:

$$\begin{aligned} &\psi_i(\mathbf{10}), \phi_i(\mathbf{5}^*), \\ &H(\mathbf{5}), \bar{H}(\mathbf{5}^*), \theta(\mathbf{50}), \bar{\theta}(\mathbf{50}^*), \Sigma(\mathbf{75}), \end{aligned}$$

where $i(= 1-3)$ represents family index. This original MD model is incomplete, since the SU(5) gauge coupling constant $\alpha_5(\equiv g_5^2/4\pi)$ blows up below the gravitational scale $M_G \simeq 2.4 \times 10^{18} \text{GeV}$ because of the large representations for $\theta, \bar{\theta}$, and Σ . One may avoid this unwanted situation by taking a mass for θ and $\bar{\theta}$ at the gravitational scale. However, in this case the colored Higgs mass M_{H_c} becomes lower than $\sim 10^{15} \text{GeV}$ which leads to too rapid nucleon decays by dimension-five operators [10].

The MMD SU(5) model [2] solves the above problem by imposing an additional

¹ Unfortunately, the MMD SU(5) model can not be excluded even if the nucleon decays are not observed by the SuperKamiokande experiments.

chiral (Peccei-Quinn) symmetry. To incorporate the Peccei-Quinn symmetry in the original MD model, we introduce a new set of chiral supermultiplets,

$$H'(\mathbf{5}), \overline{H}'(\mathbf{5}^*), \theta'(\mathbf{50}), \overline{\theta}'(\mathbf{50}^*).$$

We, then, assume the $U(1)_{PQ}$ charges for each multiplets as

$$\begin{aligned} \psi_i(\mathbf{10}) &\rightarrow e^{i\alpha/2} \psi_i(\mathbf{10}), & \phi_i(\mathbf{5}^*) &\rightarrow e^{i\beta/2} \phi_i(\mathbf{5}^*), \\ H(\mathbf{5}) &\rightarrow e^{-i\alpha} H(\mathbf{5}), & \overline{H}(\mathbf{5}^*) &\rightarrow e^{-i\frac{\alpha+\beta}{2}} \overline{H}(\mathbf{5}^*), \\ H'(\mathbf{5}) &\rightarrow e^{i\frac{\alpha+\beta}{2}} H'(\mathbf{5}), & \overline{H}'(\mathbf{5}^*) &\rightarrow e^{i\alpha} \overline{H}'(\mathbf{5}^*), \\ \theta(\mathbf{50}) &\rightarrow e^{i\alpha} \theta(\mathbf{50}), & \overline{\theta}(\mathbf{50}^*) &\rightarrow e^{i\frac{\alpha+\beta}{2}} \overline{\theta}(\mathbf{50}^*), \\ \theta'(\mathbf{50}) &\rightarrow e^{-i\frac{\alpha+\beta}{2}} \theta'(\mathbf{50}), & \overline{\theta}'(\mathbf{50}^*) &\rightarrow e^{-i\alpha} \overline{\theta}'(\mathbf{50}^*), \\ \Sigma(\mathbf{75}) &\rightarrow \Sigma(\mathbf{75}), \end{aligned} \tag{1}$$

with $3\alpha + \beta \neq 0$.

We take masses of $\theta, \overline{\theta}'$, and $\theta', \overline{\theta}$ at the gravitational scale M_G to maintain the perturbative description of the SUSY GUT. Then, we have only two pairs of Higgs multiplets, $H, \overline{H}'$, and $H', \overline{H}$, and one Higgs Σ in addition to three families of quark and lepton multiplets below the scale M_G .

The 75-dimensional Higgs Σ has the following vacuum expectation value causing the desired $SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$ breaking,

$$\begin{aligned} \langle \Sigma \rangle_{(\gamma\delta)}^{(\alpha\beta)} &= \frac{1}{2} \left\{ \delta_\gamma^\alpha \delta_\delta^\beta - \delta_\delta^\alpha \delta_\gamma^\beta \right\} V_\Sigma, \\ \langle \Sigma \rangle_{(cd)}^{(ab)} &= \frac{3}{2} \left\{ \delta_c^a \delta_d^b - \delta_d^a \delta_c^b \right\} V_\Sigma, \\ \langle \Sigma \rangle_{(b\beta)}^{(a\alpha)} &= -\frac{1}{2} \left\{ \delta_b^a \delta_\beta^\alpha \right\} V_\Sigma, \end{aligned} \tag{2}$$

where $\alpha, \beta, \dots$ are the $SU(3)_C$ indices and $a, b, \dots$ the $SU(2)_L$ indices. Integration of the heavy fields, $\theta, \overline{\theta}'$, and $\theta', \overline{\theta}$, generates Peccei-Quinn invariant masses of the colored Higgs multiplets as

$$M_{H_c} H_c^\alpha \overline{H}'_{c\alpha} + M_{\overline{H}_c} H'_c{}^\alpha \overline{H}_{c\alpha} \tag{3}$$

with

$$M_{H_c} \simeq 48 G_H G'_H \frac{V_\Sigma^2}{M_G}, \quad M_{\overline{H}_c} \simeq 48 G_{\overline{H}} G'_H \frac{V_\Sigma^2}{M_G}. \tag{4}$$

Here, the coupling constants G_H , $G_{\overline{H}}$, G'_H , and $G'_{\overline{H}}$ are expected to be $O(1)$ (see Ref. [2] for their definition). Notice that dimension-five operators for nucleon decays are completely suppressed as long as the Peccei-Quinn symmetry is unbroken.

The two pairs of Higgs doublets, H_f , $\overline{H}_f$, and H'_f , $\overline{H}'_f$, on the other hand, remain massless. They acquire masses through the Peccei-Quinn symmetry breaking. As shown in Ref. [2] we choose $U(1)_{PQ}$ charges for the Peccei-Quinn symmetry breaking fields so that only one pair of Higgs doublets H'_f and $\overline{H}'_f$ has a mass $M_{H'_f}$ of the order of the Peccei-Quinn scale $\sim (10^{10} - 10^{12})\text{GeV}$. Thus, the model is nothing but the SUSY standard model below the Peccei-Quinn scale. The Peccei-Quinn symmetry breaking generates the dimension-five operators which are, however, proportional to $M_{H'_f}/(M_{H_c}M_{\overline{H}_c})$.

We are now ready to discuss renormalization group (RG) equations for the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge coupling constants α_3 , α_2 , and α_1 in the standard model. It has been pointed out in Ref. [11] that the GUT scale and the colored Higgs mass are determined independently by using the gauge coupling constants at the electroweak scale. For the case of the MMD model we have [2] at the one-loop level

$$(3\alpha_2^{-1} - 2\alpha_3^{-1} - \alpha_1^{-1})(m_Z) = \frac{1}{2\pi} \left\{ \frac{12}{5} \ln \frac{M_{H_c}^{eff}}{m_Z} - 2 \ln \frac{m_{SUSY}}{m_Z} - \frac{12}{5} \ln(1.7 \times 10^4) \right\}, \quad (5)$$

$$(5\alpha_1^{-1} - 3\alpha_2^{-1} - 2\alpha_3^{-1})(m_Z) = \frac{1}{2\pi} \left\{ 12 \ln \frac{M_V^2 M_\Sigma}{m_Z^3} + 8 \ln \frac{m_{SUSY}}{m_Z} + 36 \ln(1.4) \right\}. \quad (6)$$

Here, the effective mass for the colored Higgs $M_{H_c}^{eff} = M_{H_c}M_{\overline{H}_c}/M_{H'_f}$. Notice that the last values in Eqs. (5) and (6) come from the mass splitting among the components in $\Sigma(\mathbf{75})$. M_V is the mass for the GUT gauge bosons ($M_V = 2\sqrt{6}g_5 V_\Sigma$),² and M_Σ the mass of the heaviest component of $\Sigma(\mathbf{75})$ ($M_\Sigma = (10/3)\lambda_{75} V_\Sigma$). The definition of Yukawa coupling constant λ_{75} is given in Ref. [2].

² We find a mistake in a form of M_V in Ref. [2].

To perform a quantitative analysis we use two-loop RG equations between the electroweak and the GUT scale. For GUT scale threshold corrections we use the one-loop result. We use the mass spectrum derived from the minimum supergravity instead of the common mass m_{SUSY} for superparticles to calculate one-loop threshold corrections at the SUSY-breaking scale. Using the experimental data $\alpha_{em}^{-1}(m_Z) = 127.90 \pm 0.09$, $\sin^2 \theta_W(m_Z) = 0.2314 \pm 0.0004$ [12] and $\alpha_3(m_Z) = 0.118 \pm 0.004$ [7], we obtain the GUT scale as

$$9.5 \times 10^{15} \text{ GeV} \leq (M_V^2 M_\Sigma)^{1/3} \leq 2.3 \times 10^{16} \text{ GeV}. \quad (7)$$

The dimension-six operators for nucleon decays depend on M_V , and hence Eq. (7) is not sufficient to calculate the decay rate. Fortunately, the lower bound on M_V is given by the following consideration. The upper bound M_Σ is determined by requiring that the Yukawa coupling λ_{75} never blows up below the gravitational scale. By solving the one-loop RG equation for λ_{75} we find³

$$M_\Sigma \leq 1.3 M_V, \quad (8)$$

which leads to together with Eq. (7)

$$M_V \geq 8.7 \times 10^{15} \text{ GeV}. \quad (9)$$

This gives the lower bound for the proton life time as

$$\begin{aligned} \tau(p \rightarrow e^+ \pi^0) &= 2.9 \times 10^{35} \times \left(\frac{M_V}{10^{16} \text{ GeV}} \right)^4 \left(\frac{0.0058 \text{ GeV}^3}{\alpha} \right)^2 \text{ years} \\ &\geq 1.6 \times 10^{35} \text{ years}. \end{aligned} \quad (10)$$

Here, α is the hadron matrix element⁴ and we have used a lattice value $\alpha = 0.0058 \text{ GeV}^3$ [13].

One may consider that threshold corrections from some additional particles at the GUT scale could decrease the GUT scale. However, if one introduces additional multiplets at the GUT scale, the SU(5) gauge coupling constant tends to blow up

³ When λ_{75} is much larger than g_5 at the gravitational scale, it rapidly closes to the quasi-infrared fixed point at the GUT scale, and on the fixed point $M_\Sigma/M_V = 1.28$.

⁴See Ref. [10] for the definition of α .

below the gravitational scale. Simple choices may be pairs of $\mathbf{5} + \mathbf{5}^*$ and/or pairs of $\mathbf{10} + \mathbf{10}^*$. We easily see from Eq. (6) that the pairs of $\mathbf{5} + \mathbf{5}^*$ do not change the GUT scale $(M_V^2 M_\Sigma)^{1/3}$. As for $\mathbf{10} + \mathbf{10}^*$ we have checked that any significant change of the GUT scale is not obtained by their threshold corrections for a natural range of mass parameters.

We are now led to consider nonrenormalizable interactions suppressed by $1/M_G$. The most dominant operator contributing to the gauge coupling constants α_3 , α_2 , and α_1 is

$$\int d^2\theta \frac{\delta}{M_G} \mathcal{W}_B^A \cdot \mathcal{W}_D^C \Sigma_{(AC)}^{(BD)} + \text{h.c.}, \quad (11)$$

where $A, B, \dots$ are SU(5) indices. Here, $\mathcal{W}$ is the SU(5) field-strength superfields whose kinetic term is given by

$$\int d^2\theta \frac{1}{8g_5^2} \mathcal{W}_B^A \cdot \mathcal{W}_A^B + \text{h.c.} \quad (12)$$

The above nonrenormalizable term yields corrections to $\alpha_i (i = 1 - 3)$ at the GUT scale as

$$\begin{aligned} \alpha_3^{-1} &= \alpha_5^{-1} - 4 \left(4\pi\delta \frac{V_\Sigma}{M_G} \right), \\ \alpha_2^{-1} &= \alpha_5^{-1} - 12 \left(4\pi\delta \frac{V_\Sigma}{M_G} \right), \\ \alpha_1^{-1} &= \alpha_5^{-1} + 20 \left(4\pi\delta \frac{V_\Sigma}{M_G} \right). \end{aligned} \quad (13)$$

Then, Eqs. (5) and (6) are modified as

$$\begin{aligned} (3\alpha_2^{-1} - 2\alpha_3^{-1} - \alpha_1^{-1})(m_Z) &= \frac{1}{2\pi} \left\{ \frac{12}{5} \ln \frac{M_{H_c}^{eff}}{m_Z} - 2 \ln \frac{m_{\text{SUSY}}}{m_Z} \right. \\ &\quad \left. - 48 \left(8\pi^2\delta \frac{V_\Sigma}{M_G} \right) - \frac{12}{5} \ln(1.7 \times 10^4) \right\}, \end{aligned} \quad (14)$$

$$\begin{aligned} (5\alpha_1^{-1} - 3\alpha_2^{-1} - 2\alpha_3^{-1})(m_Z) &= \frac{1}{2\pi} \left\{ 12 \ln \frac{M_V^2 M_\Sigma}{m_Z^3} + 8 \ln \frac{m_{\text{SUSY}}}{m_Z} \right. \\ &\quad \left. + 144 \left(8\pi^2\delta \frac{V_\Sigma}{M_G} \right) + 36 \ln(1.4) \right\}. \end{aligned} \quad (15)$$

We see that the GUT scale $(M_V^2 M_\Sigma)^{1/3}$ decreases easily by a factor ~ 3 for $\delta V_\Sigma/M_G = 0.0035$.⁵ As a consequence the effective mass for the colored Higgs multiplets $M_{H_c}^{eff}$

⁵ Introduction of a similar nonrenormalizable operator to Eq. (11) suppressed by $1/M_G$ does not change the GUT scale in the minimal SUSY SU(5) GUT [14].

increases by almost 240 times compared with the values for $\delta = 0$. This is rather welcome to suppress the dimension-five operators for nucleon decays since they are proportional to $1/M_{H_c}^{eff}$. Notice that the presence of the nonrenormalizable operator in Eq. (11) lowers further the value of $\alpha_3(m_Z)$ for $\delta > 0$. This over reduction of $\alpha_3(m_Z)$ is, however, compensated by raising the $M_{H_c}^{eff}$.

In conclusion, we have shown in this letter that introduction of the nonrenormalizable interaction may decrease the GUT scale by a factor ~ 3 in the MMD SU(5) model [2]. In Fig. 1, we show the theoretical lower bounds on the proton lifetime $\tau(p \rightarrow e^+\pi^0)$. We see that they lie within the observable range at SuperKamiokande for a reasonable parameter region $\delta \sim O(1)$. A recent report [15] from SuperKamiokande shows that one event of the mode $e^+\pi^0$ still survives the cut criterion for backgrounds. If this is indeed a real signal for $p \rightarrow e^+\pi^0$ decay, we may be led to a serious consideration of physics at the gravitational scale.




Figure 1: The theoretical lower bounds on the proton lifetime $\tau(p \rightarrow e^+ \pi^0)$ as a function of the parameter $(\delta V_\Sigma / M_G) \times 10^2$. Here, we have assumed a lattice value of the hadron matrix element $\alpha = 0.0058 \text{ GeV}^3$ [13]. The shaded area is excluded by the recent SuperKamiokande experiments ($\tau(p \rightarrow e^+ \pi^0) > 7.9 \times 10^{32} \text{ years}$ [15]).

References

- [1] A. Masiero, D.V. Nanopoulos, K. Tamvakis, and T. Yanagida, *Phys. Lett.* **B115** (1982) 380;
B.Grinstein, *Nucl. Phys.* **B206** (1982) 387.
- [2] J. Hisano, T. Moroi, K. Tobe, and T. Yanagida, *Phys. Lett.* **B342** (1995) 138.
- [3] S.M. Barr, *Phys. Lett.* **B112** (1982) 219; *Phys. Rev.* **D40** (1989) 2457;
J.P. Derendinger, J.E. Kim, and D.V. Nanopoulos, *Phys. Lett.* **B139** (1984) 170.
- [4] S. Dimopoulos and F. Wilczek, University of California at Santa Barbara, Institute for Theoretical Physics Report No. NSF-ITP-82-07;
M. Srednicki, *Nucl. Phys.* **B202** (1982) 327.
- [5] K. Inoue, A. Kakuto, and H. Takano, *Prog. Theor. Phys.* **75** (1986) 664;
A.A. Anselm and A.A. Johansen, *Phys. Lett.* **B200** (1988) 331.
- [6] T. Yanagida, *Phys. Lett.* **B344** (1995) 211;
T. Hotta, K.I. Izawa, and T. Yanagida, *Phys. Rev.* **D53** (1996) 3913.
- [7] W.J. Stirling, Talk given at “New Nonperturbative Methods and Quantization of the Light Cone”, France, Mar. 1997 (hep-ph/9709429).
- [8] J. Bagger, K. Matchev, and D. Pierce, *Phys. Lett.* **B348** (1995) 443;
B. Brahmachari and R.N. Mohapatra, *Int. J. Mod. Phys.* **A11** (1996) 1699;
A. Dedes, A.B. Lahanas, J. Rizos, and K. Tamvakis, *Phys. Rev.* **D55** (1997) 2955.
- [9] N. Sakai and T. Yanagida, *Nucl. Phys.* **B197** (1982) 533;
S. Weinberg, *Phys. Rev.* **D26** (1982) 287.
- [10] J. Hisano, H. Murayama, and T. Yanagida, *Nucl. Phys.* **B402** (1993) 46.
- [11] J. Hisano, H. Murayama, and T. Yanagida, *Phys. Rev. Lett.* **69** (1992) 1014.
- [12] Particle Data Group, *Phys. Rev.* **D54** (1996) 1.
- [13] M.B. Gavela, S.F. King, C.T. Sachrajda, G. Martinelli, M.L. Paciello, and B. Taglienti, *Nucl. Phys.* **B312** (1989) 269.

- [14] L.J. Hall and U. Sarid, *Phys. Rev. Lett.* **70** (1993) 2673.
- [15] SuperKamiokande Collaboration, Talk Given at the meeting of the Physical Society of Japan, Tokyo, Sept. 1997.